Name: Thomawin Pattamaphol (Win)

bet=a=8, b=11, c=3

$$3x_1 - 40c_2 + 8x_3 = -1$$

$$-3x_1 + 11x_2 + 3x_3 = 1$$

$$8x_1 + 8x_2 - 11x_3 = 32$$

Augmented matrix

$$\begin{bmatrix}
3 & -4 & 8 & | & -1 \\
-3 & 11 & 3 & | & 1 \\
8 & 8 & -11 & | & 32
\end{bmatrix}
\xrightarrow{R2 + R1 \rightarrow R2}
\xrightarrow{R3 - (\frac{8}{3})R1 \rightarrow R3}
\begin{bmatrix}
3 & -4 & 8 & | & -1 \\
0 & 7 & 11 & | & 0 \\
0 & \frac{56}{3} & -\frac{97}{3} & | & \frac{104}{3}
\end{bmatrix}
\xrightarrow{R3 - \frac{9}{3}R2 \rightarrow R3}
\begin{bmatrix}
3 & -4 & 8 & | & -1 \\
0 & 7 & 11 & | & 0 \\
0 & 0 & -185 & | & 104
\end{bmatrix}$$

Baile substitution.

$$\frac{-183\alpha_3}{-183} = \frac{104}{-183}$$

$$\frac{-183}{2} = \frac{104}{183}$$

$$72_{2}+112_{3}=0$$

$$72_{2}+112_{3}=0$$

$$2=-\frac{11}{7}\cdot\left(-\frac{104}{185}\right)$$

$$=\frac{11\cdot104}{7\cdot185}$$

$$=\frac{1144}{1095}$$

$$3x_1 - 4x_2 + 8x_3 = -1$$

$$3x_1 = -1 + 4x_2 - 8x_3$$

$$2 - 1 + 4\left(\frac{1144}{1295}\right) - 8\left(\frac{104}{183}\right)$$

$$= \frac{607}{259}$$

Geometrie interpretation.

The Three equations represent three different planes, in which, they me not parallel & not concurrent along a line, i.e. threy intersect at a point.

And the point is (607, 1144, -104)

## Name: Thanawin Pattanaphol - LLM used: ChatGPT

Given system:

$$\begin{cases} 3x_1 - 4x_2 + ax_3 = -1, \\ -3x_1 + bx_2 + cx_3 = 1, \\ ax_1 + 8x_2 - bx_3 = 32, \end{cases}$$
 where  $a = 8, b = 11, c = 3.$ 

$$\Rightarrow \begin{cases} 3x_1 - 4x_2 + 8x_3 = -1, \\ -3x_1 + 11x_2 + 3x_3 = 1, \\ 8x_1 + 8x_2 - 11x_3 = 32. \end{cases}$$

$$\Rightarrow \begin{cases} 3x_1 - 4x_2 + 8x_3 = -1 \\ -3x_1 + 11x_2 + 3x_3 = 1 \\ 8x_1 + 8x_2 - 11x_3 = 32 \end{cases}$$

Turn into augmented matrix:

Augmented matrix: 
$$\begin{bmatrix} 3 & -4 & 8 & | & -1 \\ -3 & 11 & 3 & | & 1 \\ 8 & 8 & -11 & | & 32 \end{bmatrix}$$

## Gauss elimination (by row operations)

Start with the augmented matrix and swap R1↔R3R\_1\leftrightarrow R\_3R1↔R3 to use an 8-pivot:

$$R_1 \leftrightarrow R_3 \Rightarrow \begin{bmatrix} 8 & 8 & -11 & | & 32 \\ -3 & 11 & 3 & | & 1 \\ 3 & -4 & 8 & | & -1 \end{bmatrix}$$

Eliminate x1 from R2, R3

$$R_2 \leftarrow 8R_2 + 3R_1, \quad R_3 \leftarrow 8R_3 - 3R_1$$

$$\Rightarrow \begin{bmatrix} 8 & 8 & -11 & | & 32 \\ 0 & 112 & -9 & | & 104 \\ 0 & -56 & 97 & | & -104 \end{bmatrix}$$

## Name: Thanawin Pattanaphol - LLM used: ChatGPT

Eliminate x3 from R3:

$$R_3 \leftarrow 2R_3 + R_2 \Rightarrow \begin{bmatrix} 8 & 8 & -11 & | & 32 \\ 0 & 112 & -9 & | & 104 \\ 0 & 0 & 185 & | & -104 \end{bmatrix}$$

## **Back Substitution**

Back substitution: 
$$\begin{cases} x_3 = \frac{-104}{185} \\ 112x_2 - 9x_3 = 104 \Rightarrow x_2 = \frac{1144}{1295} \\ 8x_1 + 8x_2 - 11x_3 = 32 \Rightarrow x_1 = \frac{607}{259} \end{cases}$$

So

$$x_1 = \frac{607}{259}, \quad x_2 = \frac{1144}{1295}, \quad x_3 = -\frac{104}{185}$$